

Applications of Virtual Environments in Medicine*

G. Riva

Applied Technology for Neuro-Psychology Lab., Istituto Auxologico Italiano, Milan, Italy

Summary

Objective: This paper intends to investigate the role of virtual reality (VR) in medicine. In particular it outlines the current state of research and technology that is relevant to the development of effective virtual environments in medicine.

Method: After describing the two different visions of VR we can find in medicine – the presentation of virtual objects to all of the human senses in a way identical to their natural counterpart, and a new human-computer interaction paradigm in which users are active participants within a computer-generated three-dimensional virtual world – the paper presents some of the most interesting applications actually developed in the area. Finally, it discusses the clinical principles, technological devices and safety issues associated with the use of VR in medicine.

Conclusions: The possible impact of VR on health care could be even higher than the one offered by the new communication technologies like Internet. In fact, VR is at the same time technology, a communication interface and an experience: a communication interface based on interactive 3D visualization, able to collect and integrate in single real-like experience different inputs and data sets. However, significant efforts are still required to move VR into commercial success and therefore routine clinical use.

Keywords

Virtual reality, medical education, surgical simulation, neuropsychological rehabilitation

Methods Inf Med 2003; 42: 524–34

1. Introduction

As recently noted by Satava and Jones [1], the advantages of virtual environments (VEs) to health care can be summarized in a single word: revolutionary. Since the development of methods of electronic communication clinicians have been using information and communication technologies in health care: telegraphy, telephony, radio and television have been used for distance medicine since mid 19th century [2]. However, rapid and far-reaching technological advances are changing the ways in which people relate, communicate, and live. Technologies that were hardly used ten years ago, such as the Internet, e-mail, and video teleconferencing are becoming familiar methods for diagnosis, therapy, education and training. However, the possible impact of virtual reality (VR) on health care is even higher than the one offered by the new communication technologies [3]. In fact, VR is a technology, a communication interface and an experience [4]. This is why the research in the virtual reality field is moving fast. If we check the two leading clinical databases – MEDLINE and PSYCINFO – using the “virtual reality” keyword we can find 951 papers listed in MEDLINE and 708 in PSYCINFO (all fields query, accessed June 9, 2003).

From the analysis of the retrieved papers we can find that the first health care applications of VR started in the early '90s by the need of medical staff to visualize complex medical data, particularly during surgery and for surgery planning [5]. Actually, surgery-related applications of VR fall

mainly into three classes: surgery training, surgery planning and augmented reality for surgery sessions in open surgery, endoscopy, and radiosurgery. A couple of years later, the scope of VR applications in medicine has broadened to include neuropsychological assessment and rehabilitation [6, 7].

In recent years, VR has generated both great excitement and great confusion. These factors are evident in the extensive material published in both scientific and popular press, and in the unrealistic expectations on the part of the health care professionals (8). In this paper we try to outline the current state of research and technology that is relevant to the development of VEs in medicine. Moreover, we discuss the clinical principles, technological devices and safety issues associated with the use of virtual reality in medicine.

2. The Role of VR in Health Care

2.1 The Two Faces of VR in Health Care

For many health care professionals VR is first of all a technology. Since 1986, when Jaron Lamier used the term for the first time, VR has been usually described as a collection of technological devices: a computer capable of interactive 3D visualization, a head-mounted display and data gloves equipped with one or more position trackers. The trackers sense the position and orientation of the user and report that information to the computer that updates (in real time) the images for display.

However, the analysis of the different VR applications clearly shows that the focus on technological devices is different

* This paper is an updated version of an invited review paper that appeared in Haux, R., Kulikowski, E. (eds.) (2003). IMIA Yearbook of Medical Informatics 2003: Quality of Health Care: The Role of Informatics, pp. 159-69, Stuttgart: Schattauer.

according to the goals of the health care provider.

For instance, Rubino et al. [9], McCloy and Stone [10], and Székely and Satava [11] in their reviews share the same vision of VR: “a collection of technologies that allow people to interact efficiently with 3D computerized databases in real time using their natural senses and skills” [10]. This definition lacks any reference to head mounted displays and instrumented clothing such as gloves or suits. In fact, less than 20% of VR health care applications in medicine are actually using any immersive equipment.

However, if we shift our attention on behavioral sciences, where immersive devices are used by more than 50% of the applications, VR is described as “an advanced form of human-computer interface that allows the user to interact with and become immersed in a computer-generated environment in a naturalistic fashion” [12]. In fact, to achieve the feeling of “being there” the VR applications use ■ specialized devices as head-mounted displays, tracking systems, earphones, gloves, and sometimes haptic-feedback devices.

These two definitions underline two different visions of VR. For physicians and surgeons, the ultimate goal of VR is the presentation of virtual objects to all of the human senses in a way identical to their natural counterpart [11]. As noted by Satava and Jones [1], as more and more of the medical technologies become information-based, it will be possible to represent a patient with higher fidelity to a point that the image may become a surrogate for the patient – the *medical avatar*. In this sense, an effective VR system should offer real-like body parts or avatars that interact with external devices such as surgical instruments as near as possible to their real models.

For clinical psychologists and rehabilitation specialists the ultimate goal is radically different [13, 14]. They use VR to provide a new human-computer interaction paradigm in which users are no longer simply external observers of images on a computer screen but are active participants within a computer-generated three-dimensional virtual world. Within the VE the patient has the possibility of learning to manage a problematic situation related to his/her dis-

turbance. The key characteristics of virtual environments for these professionals are both the high level of control of the interaction with the tool without the constraints usually found in computer systems, and the enriched experience provided to the patient [12]. Virtual environments are highly flexible and programmable. They enable the therapist to present a wide variety of controlled stimuli, such as a fearful situation, and to measure and monitor a wide variety of responses made by the user. This flexibility can be used to provide systematic restorative training that optimize the degree of transfer of training or generalization of learning to the person’s real world environment [15].

Moreover, virtual reality systems open the input channel to the full range of human gestures: in rehabilitation it is possible to monitor movements or actions from any body part or many body parts at the same time. On the other side, with disabled patients feedbacks and prompts can be translated into alternate and/or multiple senses [16].

2.2 VR as Communication Interface

As we have just seen, if we consider VR mainly as a technology we have two different visions of VR related to the final goal of the health care professional. But what these two visions have in common?

The starting point for answering to this question is a definition of VR presented by Heim. According to this author [17], VR is “an immersive, interactive system based on computable information... an experience that describes many life activities in the information age” (p. 6). In particular he describes the VR experience around its “three I’s”: immersion, interactivity and information intensity. Developing this position, Bricken [18] identifies the core characteristic of VR in the inclusive relationship between the participant and the virtual environment, where direct experience of the immersive environment constitutes communication. According to this position, VR can be considered as the leading edge of a general evolution of present communication interfaces like television, computer

and telephone [19, 20]. The main characteristic of this evolution is the full immersion of the human sensorimotor channels into a vivid and global communication experience [21].

Following this approach, it is also possible to define VR in terms of human experience [22] “a real or simulated environment in which a perceiver experiences telepresence”, where telepresence can be described as the “experience of presence in an environment by means of a communication medium” (pp. 78-80).

This position better clarifies the possible role of VR in medicine: a communication interface based on interactive 3D visualization, able to collect and integrate different inputs and data sets in a single real-like experience. It is up to the health care provider to decide if the VR application will be more focused on the integration of different data sets or on the realism of the virtual experience. Considering VR as a communication interface also helps health care developers to focus their efforts.

Most of the work in this area is trying to improve the efficacy of a VE by providing to the user a more “realistic” experience, such as adding physical qualities to virtual objects or improving the graphical resolution. But is it really so important for the effectiveness of a medical VE this focus on the graphical characteristics?

Probably, apart from some high-end surgical applications, the answer is no. More than the richness of available images, the efficacy of a virtual environment depends on the level of interaction/interactivity which actors have in both “real” and simulated environments [23]. According to Sastri and Boyd [23] a VE, particularly when it is used for real world applications, is effective when “the user is able to navigate, select, pick, move and manipulate an object much more naturally” (pp. 235). In this sense, emphasis shifts from quality of image to freedom of interaction, from the graphic perfection of the system to the affordances provided to the users in the environment [24]. Further, as the underlying enabling technologies continue to evolve and allow us to design more useful and usable structural virtual environments the next important challenge will involve populating these

environments with virtual representations of humans (avatars) [25].

This is possible because the key characteristic of VR, differentiating it from other media or communication systems, is the sense of *presence* [26, 27]. What is presence? Even if usually presence is defined as the “sense of being there” [22], or as the “feeling of being in a world that exists outside of the self” [28], it is now widely acknowledged that presence can be considered as a neuropsychological phenomenon [20, 26, 29-33]. In particular, Riva and Waterworth described presence as a defining feature of self, related to the evolution of a key feature of any central nervous system [28]: the embedding of sensory-referred properties into an internal functional space. More in particular, without the emergence of the sense of presence it is impossible for the nervous system to separate between an external world and the internal one. If in simple organisms, this separation involved only a correct coupling between perceptions and movements, in humans it also requires the shift from meaning-as-comprehensibility to meaning-as-significance. Meaning-as-comprehensibility refers to the extent to which the event fits with our view of the world (for example, as just, controllable, and nonrandom) whereas meaning-as-significance refers to the value or worth of the event for us [34]. Following this point, contributions to the intensity of the sense of presence come from three layers of the self recently defined by Damasio [35]: proto self, core self and autobiographical self. The more the three layers are integrated (focused on the same events) the stronger the intensity of the presence feeling [28]. This means that having two equally stimulating virtual environments, humans are more present in the one more relevant to their own goals.

This approach has recently received the status of international standard, through the International Organization for Standardization’s ISO 13407 “Human centered design for interactive systems”. According to the ISO 13407 standard [36], human-centered design requires:

- the active involvement of users;
- clear understanding of use and task requirements;

- appropriate allocation of function;
- the iteration of design solutions;
- a multi-disciplinary design team;

and it is based around the following processes:

- understand and specify the context of use;
- specify the user and organizational requirements;
- produce designs and prototypes;
- carry out user-based assessment.

A sample of VE developed using the ISO 13407 guidelines is the IERAPSI surgical training system [10, 37].

3. Applications of Virtual Reality in Medicine

3.1 Medical Education

The teaching of anatomy is mainly illustrative, and the application of VR to such teaching has great potential [38]. Through 3-D visualization of massive volumes of information and databases, clinicians and students can understand important physiological principles or basic anatomy [39]. For instance, VR can be used to explore the organs by “flying” around, behind, or even inside them. In this sense VEs can be used both as didactic and experiential educational tools, allowing a deeper understanding of the interrelationship of anatomical structures that cannot be achieved by any other means, including cadaveric dissection.

A significant step towards the creation of VR anatomy textbooks was the acquisition of the Visible Human male and female data made in August of 1991 by the University of Colorado School of Medicine [40]. The Visible Human female data set contains 5189 digital anatomical images obtained at 0.33-mm intervals (39 Gbyte). The male data set contains 1971 digital axial anatomical images obtained at 1.0-mm intervals (15 Gbyte) [41]. Actually, the US National Library of Medicine in partnership with other US government research agencies has begun the development of a tool

kit of computational programs capable of automatically performing many of the basic data handling functions required for using Visible Human data in applications [42].

The National Library of Medicine made the data sets available under a no-cost license agreement over the Internet. And this allowed the creation of a huge number of educational VEs. In their recent edited book Westwood and colleagues [43] report more than ten different educational and visualization applications.

In the future we can expect the development of different VR dynamic models illustrating how various organs and systems move during normal or diseased states, or how they respond to various externally applied forces (e.g., the touch of a scalpel).

Apart from anatomical training, VR has been used for teaching the skill of performing a 12-lead ECG [44]. In all these cases, VR simulators allowed the acquisition of necessary technical skills required for the procedure.

3.2 Surgical Simulation and Planning

Surgeons know well that in training there is no alternative to hands-on practice. However, students wishing to learn laparoscopic procedures face a tough path [45]: usually they start using laparoscopic cholecystectomy trainers consisting of a black box in which endoscopic instruments are passed through rubber gaskets. After, the students begin practicing these techniques on inanimate tissues, when allowed by their cost and availability. Obviously, there is a substantial difference for students between training with artificial or inanimate tissues and supervised procedures on real patients. This is why in early 1990s different research teams tried to develop VE simulators [46, 47]. The science of virtual reality provides an entirely new opportunity in the area of simulation of surgical skills using computers for training, evaluation, and eventually certification [48]. However the first simulators were limited by low-resolution graphics, the lack of tactile input and force feedback and the lack of realistic deformation of organs. In the last years a new generation

of simulator has appeared that has shown improved training efficacy over traditional methods [49, 50, Schijven, 2003 #1455]. For instance, a randomized trial using the Minimally Invasive Surgery Training-Virtual Reality (MIST-VR) trainer [51, 52] showed that VR simulation was effective in training the novice to perform basic laparoscopic skills (see Fig. 1).

Another typical use of visualization applications is the planning of surgical and neuro-surgical procedures [53-55]. The planning of these procedures usually relies on the studies of series of two-dimensional MR (Magnetic Resonance) and/or CT (Computer Tomography) images, which have to be mentally integrated by surgeons into a three-dimensional concept. This mental transformation is difficult, since complex anatomy is represented in different scanning modalities, on separate image series, usually found in different sites/departments. A VR-based system is capable of incorporating different scanning modalities coming from different sites providing a simple to use interactive three-dimensional view. Within the Virtual Collaborative Clinic project, NASA researchers developed Cyberscalpel, a typical VR-based surgical system for planning and practice [56]. To plan the operation of a patient with a cancer of the jaw, the upper and lower jaws were reconstructed using Cyberscalpel starting from a CT scan. The scan was reduced to 20,000 polygons and the final model used to prove how fibular bone could be sectioned to mimic and replace the jaw pieces.

Finally, the increased pressure to reduce the use of animals in technical training has led to use VR in teaching microsurgery [57]. This new technology may prove to be a cost-effective, portable, and nonhazardous way forward in microsurgical training.

3.3 Virtual Endoscopy

Every year the screening for cancer requires the performance of over 2 million video colonoscopic procedures. However, these procedures are not perfect:

- all endoscopic procedures are invasive;
- the patients are subject to complications such as perforation, bleeding, etc.



Fig. 1 Minimally Invasive Surgery Training-Virtual Reality (MIST-VR) trainer (Mentice Medical Simulation AB, Gothenburg, Sweden)

- the cost for a typical colonoscopy is significant.

To overcome these problems, different researchers are investigating the possibility of virtual endoscopy [9, 58]. Virtual endoscopy is a new procedure that fuses computed tomography with advanced techniques for rendering three-dimensional images to produce views of the organ similar to those obtained during “real” endoscopy. A virtual endoscopy is performed by using a standard CT scan or MRI scan [1], reconstructing the organ of interest into a 3D model, and then performing a fly through it. Typical examples include the colon, stomach, esophagus, tracheo-bronchial tree (bronchoscopy), sinus bladder, ureter and kidneys (cystoscopy), pancreas or biliary tree [59].

Virtual endoscopy is completely non-invasive and thus without known complications [60]. The actual cost is less of traditional endoscopy, since it is performed in the same place and manner as all imaging modalities, utilizes the same staff, and has no consumable materials.

3.4 VR in Neuro-Psychological Assessment and Rehabilitation

VR is starting to play an important role in clinical psychology [61, 62], that is expected to increase in the next years. According to a recent positioning paper on the future of psychotherapy [63], the use of VR and computerized therapies are ranked respectively 3rd and 5th out of 38 psychotherapy interventions that are predicted to increase in the next 10 years.

In most VEs for clinical psychology VR is used to simulate the real world and to assure the researcher full control of all the parameters implied. VR constitutes a highly flexible tool, which makes it possible to program an enormous variety of procedures of intervention on psychological distress. The possibility of structuring a large amount of controlled stimuli and, simultaneously, of monitoring the possible responses generated by the user of the virtual world offers a considerable increase in the likelihood of therapeutic effectiveness, as compared to traditional procedures [20]. In particular, a key advantage offered by VR is the possibility for the patient to manage successfully a problematic situation related to his/her disturbance. Using VR in this way, the patient is more likely not only to gain an awareness of his/her need to do something to create change but also to experience a greater sense of personal efficacy.

In general, these techniques are used as triggers for a broader empowerment process. In psychological literature *empowerment* is considered a multi-faceted construct reflecting the different dimensions of being psychologically enabled, and is conceived of as a positive additive function of the following three dimensions [64]:

- *perceived control*: includes beliefs about authority, decision-making skills, availability of resources, autonomy in the scheduling and performance of work, etc;
- *perceived competence*: reflects role-mastery, which besides requiring the skillful accomplishment of one or more assigned tasks, also requires successful coping with non-routine role-related situations;

Table 1 VR hardware

| Workstation | Indicative Prices (as 01 Jul 03) |
|---|-------------------------------------|
| SGI Onyx350 visualization system with InfinitePerformance & InfiniteReality4 graphics | 250000 US\$ |
| SGI Octane2, V12 Graphic Card, 2x400MHz processors, 512 Mbyte Ram, 18 Gbyte Hard Disk | 18000 US\$ |
| Xeon branded PC, 2x3Ghz processors, 1Gbyte Ram, 2x200 Gbyte Hard Disk and 17" monitor | 4200 US\$ |
| Pentium IV or Athlon XP branded PC, 3 Ghz processor, 512 Mbyte Ram, 200 Gbyte Hard Disk and 17" monitor | 2600 US\$ |
| Consumer graphic cards | |
| Nvidia GeForceFX 5900 Ultra 128 Mbyte Vram AGP | 450 US\$ |
| ATI Radeon 9800 Pro All-in-Wonder 128 Mbyte Vram AGP | 400 US\$ |
| Professional graphic cards | |
| Quadro4 900XGL 128 Mbyte Vram AGP | 1200 US\$ |
| Fire GL X1 256 Mbyte Vram AGP | 1200 US\$ |
| Tracking system | |
| Polhemus Fastrak | 7000 US\$ |
| Ascension PC Flock of Birds | 2200 US\$ |
| Intersense Intertrax 2 | 1100 US\$ |
| 3D Shutter Glasses | |
| StereoEyes Wireless | 320 US\$ |
| Elsa 3D Revelator IR | 180 US\$ |
| VRex Cordless | 100 US\$ |
| Head Mounted Display | |
| Kaiser Proview XL 40/50 (XGA resolution – 3D, wide fov) | 50000 US\$ |
| N-visor Datavisor Hi-res (XGA resolution – 2D, wide fov) | 35000 US\$ |
| Daeyang I-Visor DH4400 VP 3D (SVGA resolution – 3D) | 1900 US\$ |
| Olympus Eye-Trek FMD-700 (SVGA resolution – 2D) | 1300 US\$ |
| Daeyang I-Visor DH4400 VP (SVGA resolution – 2D) | 1200 US\$ |
| Olympus Eyetrek 250 W (Video output only – 2D) | 600 US\$ |
| Sony Glasstron PLM-A35 (Video output only – 2D) | 500 US\$ |
| VR Gloves | |
| Pinch Glove | 2000 US\$ |
| 5DT Right Hand | 650 US\$ |

- *goal internalization*: this dimension captures the energizing property of a worthy cause or exciting vision provided by the organizational leadership.

Virtual reality can be considered the preferred environment for the empowerment process, since it is a special, sheltered setting where patients can start to explore and act without feeling threatened. In this sense the virtual experience is an “empowering environment” that therapy provides for patients. As noted by Botella [65], nothing the patients fear can “really” happen to them in VR. With such assurance, they can freely explore, experiment, feel, live, and experience feelings and/or thoughts. VR thus becomes a very useful intermediate step between the therapist’s office and the real world.

Even if the clinical rationale behind the use of VR is now clear, much of this research growth, however, has been in the form of feasibility studies and pilot trials. As a result there is still limited convincing evidence coming from controlled studies (see Table 2), of the clinical advantages of this approach. Up to now the clinical effectiveness of VR was verified in the treatment of these six psychological disorders: acrophobia [66-68], spider phobia [69], panic disorders with agoraphobia [70], body image disturbances [71], binge eating disorders [72, 73] (see Fig. 2), and fear of flying [74-78].

In the cognitive rehabilitation area the situation is even worse. Even if different case studies and review papers suggest the use of VR in this area [12, 15, 79-85] there are no controlled clinical trials to support this position. A better situation can be found in the assessment of cognitive functions in persons with acquired brain injuries. In this area VR assessment tools are effective and characterized by good psychometric properties [86-90]. A typical example of these applications is ARCANA. Using a standard tool (Wisconsin Card Sorting Test – WCST) of neuropsychological assessment as a model, Pugnetti and colleagues have created ARCANA: a virtual building in which the patient has to use environmental clues in the selection of appropriate choices (doorways) to navigate



Fig. 2
The Virtual Reality for Eating Disorders Modification - VREDIM (Istituto Auxologico Italiano I.R.C.C.S., Milan, Italy)

through the building. The doorway choices vary according to the categories of shape, color, and number of portholes. The patient is also required to refer to the previous doorway for clues to appropriately make his/her next choice. After the choice criteria are changed, the patient must shift the cognitive set, analyze clues, and devise a new choice strategy. The parameters of this system are fully adjustable so that training applications can follow initial standardized assessments.

4. VR Hardware and Software

For many years one of the main obstacles to the development of VR applications was the price of the equipment: a typical VR system required a costly fridge-size Silicon Graphic workstation in the range of 250,000 US\$ and up. Even if high-end applications still require powerful workstations such as SGI Onyx or Octane (see Table 1), during the last two years about 65% of the VR applications for health care were developed for being used on PC platforms.

The significant advances in PC hardware that have been made over the last five years, are transforming PC-based VR into a reality. The cost of a basic desktop VR system has gone down by many thousand dollars since that time, and the functionality has improved dramatically in terms of graphics processing power. A simple immersive VR system now may cost less than 6000 US\$ (see Table 1).

The availability of powerful PC engines based on such computing work-horses as Intel's Xeon and IBM G4/G5 processors, and the emergence of reasonably priced, Direct 3D and OpenGL-based 3D accelerator cards allow high-end PCs to process and display interactive 3D simulations in real time.

While a standard Celeron/Duron processor with as little as 128 Mbyte of RAM can provide sufficient processing power for a simple VR simulation, a fast Pentium IV/Athlon XP-based PC (2.5 Ghz or faster) with 256 Mbyte of RAM, can transport users to a convincing virtual environment,

Table 2 Controlled trials with more than 10 patients/users included in Medline/PsycInfo (all fields query, accessed June 9, 2003)

| Authors | Paper | Sample |
|---|---|--|
| Ali, M.R., Mowery, Y., Kaplan, B., DeMaria, E.J. | (2002) Training the novice in laparoscopy. <i>Surg Endosc</i> , 16 (8), 1, 1210-1216. | 27 high school students |
| Emmelkamp, P.M.G., Bruynzeel, M., Drost, L., & van der Mast, C.A.P.G. | (2001) Virtual reality treatment in acrophobia: A comparison with exposure in Vivo. <i>Cyberpsych Behav</i> , 4(3), 335-339. | 10 acrophobia patients |
| Emmelkamp, P.M.G., Krijn, M., Hulsbosch, A.M., de Vries, S., Schuemie, M.J., van der Mast, C.A.P.G. | (2002) Virtual reality treatment versus exposure in vivo: a comparative evaluation in acrophobia. <i>Behav Res Ther</i> , 40, 509-516. | 33 acrophobia patients |
| Garcia-Palacios, A., Hoffman, H., Carlin, A., Furness, T. A., 3rd, & Botella, C. | (2002). Virtual reality in the treatment of spider phobia: a controlled study. <i>Behav Res Ther</i> , 40(9), 983-993. | 23 phobics |
| Grundman, J. A., Wigton, R. S., & Nickol, D. | (2000). A controlled trial of an interactive, web-based virtual reality program for teaching physical diagnosis skills to medical students. <i>Acad Med</i> , 75(10 Suppl), S47-49. | 121 medical students |
| Hoffman, H. G., Patterson, D. R., & Carrougner, G. J. | (2000). Use of virtual reality for adjunctive treatment of adult burn pain during physical therapy: a controlled study. <i>Clin J Pain</i> , 16(3), 244-250. | 12 burn patients |
| Maltby, N., Kirsch, I., Mayers, M., & Allen, G. | (2002). Virtual Reality Exposure Therapy for the treatment of fear of flying: A controlled investigation. <i>J Consult Clin Psych</i> , 70(5), 1112-1118. | 45 phobics |
| Riva, G., Bacchetta, M., Baruffi, M., & Molinari, E. | (2001). Virtual reality-based multidimensional therapy for the treatment of body image disturbances in obesity: a controlled study. <i>Cyberpsych Behav</i> , 4(4), 511-526. | 28 obese patients |
| Riva, G., Bacchetta, M., Baruffi, M., & Molinari, E. | Virtual reality-based multidimensional therapy for the treatment of body image disturbances in binge eating disorders: A preliminary controlled study <i>IEEE Trans Inf Tech Biom</i> , 6 (3), 224-234. | 20 binge eating patients |
| Riva, G., Bacchetta, M., Cesa, G., Conti, S. & Molinari, E. | (2003) Six-month follow-up of in-patient Experiential-Cognitive Therapy for binge eating disorders. <i>Cyberpsych Behav</i> , 6(3), 251-258. | 36 binge eating patients |
| Rothbaum, B. O., Hodges, L. F., Kooper, R., Opdyke, D., & et al. | (1995). Effectiveness of computer-generated (virtual reality) graded exposure in the treatment of acrophobia. <i>Am J Psychiatry</i> , 152(4), 626-628. | 17 college students |
| Rothbaum, B. O., Hodges, L., Smith, S., Lee, J. H., & Price, L. | (2000). A controlled study of virtual reality exposure therapy for the fear of flying. <i>J Consult Clin Psychol</i> , 68(6), 1020-1026. | 49 fear of flying patients |
| Rothbaum, B. O., Hodges, L., Anderson, P.L., Price, L., & Smith, S. | (2002) Twelve-month follow-up of virtual reality and standard exposure therapies for the fear of flying. <i>J Consult Clin Psychol</i> , 70(2), 428-432. | |
| Torkington, J., Smith, S. G., Rees, B. I., & Darzi, A. | (2001). Skill transfer from virtual reality to a real laparoscopic task. <i>Surg Endosc</i> , 15(10), 1076-1079. | 30 medical students |
| Vincelli, F., Anolli, L., Bouchard, S., Wiederhold, B.K., Zurloni, V., Riva, G. | (2003) Experiential Cognitive Therapy in the Treatment of Panic Disorders with Agoraphobia: A Controlled Study. <i>Cyberpsych Behav</i> , 6(3), 312-318. | 12 panic disorders with agoraphobia patients |
| Wiederhold, B.K., Jang, D.P., Kim, S.I., & Wiederhold, M.D. | (2002). Physiological monitoring as an objective tool in virtual reality therapy. <i>Cyberpsych Behav</i> . 5(1) 77-82. | 36 fear of flying patients, 22 non-phobics |
| Wiederhold, B.K., Jang, D.P., Kim, S.I., & Wiederhold, M.D. | (2002). A controlled trial comparing physiological responses during virtual reality exposure and imaginal exposure in flight phobics. <i>IEEE Transactions on Information Technology in Biomedicine</i> , 6 (3), 218-223. | 30 fear of flying patients |

while a dual Xeon configuration (2.7 Ghz or faster) with 1 Gbyte of RAM, OpenGL acceleration and 256 Mbyte of VRAM running Windows XP Pro rivals the horsepower of a mid-level graphics workstation.

The graphics card landscape, too, is evolving quickly. In particular, two advancements are interesting for VR users: the inclusion of a VGA-to-TV converter and tuner, the Accelerated Graphics Port (AGP) and the new faster 3D chips (GeForceFX 5900 Ultra, Radeon 9800 Pro) with 128 Mbyte or more of dedicated video Ram (VRam).

- *Accelerated Graphics Port (AGP):* The accelerated graphics port is a high-speed, point-to-point connection between the system chip set and the graphics chip. AGP provides a high-speed pipeline between the graphics accelerator and the PC's system memory: using an AGP connection, a graphics chip is able to access system memory directly through the system chip set at memory-bus speeds, reducing latency and substantially increasing performance versus standard PCI-memory transfers. The graphics card gains access to system RAM to store and execute texture bitmaps, which allows more detailed textures of unlimited size while speeding 3D rendering. When textures are large, AGP can make the difference between a smooth or choppy frame rates in 3D rendering.
- *Faster 3D cards:* In VR, performance is critical. VEs gave mainstream 3D acceleration its start, and developers have been adding a sense of realistic depth to their creations for years. However, the addition of a z-axis in rendering, as opposed to simply drawing on an x, y-coordinate plane, requires more sophisticated horsepower. In addition, VR applications contain more complex objects and complex textures: bitmap renderings of detailed surfaces (bricks, sand, or transparent water) that heighten realism. To exploit this potential a fast graphics card with a lot of video Ram is a must. Happily, the new chip sets (GeForceFX 5900 Ultra and Radeon 9800 Pro) included in consumer graphics cards have 16 times more video Ram and 5 times more 3D

acceleration than the first generation of chips (GeForce and Radeon VE) for a price tag of less than 500 US\$. Also, professional graphics cards received a significant speed bump. New Open GL cards such as the Quadro 4 900XGL or the FireGLX1 offer graphics power that rival the one provided by Unix graphic workstations.

- *VGA-to-TV converter:* One welcome feature of the new graphics cards is the inclusion of a VGA-to-TV (NTSC or PAL) converter and TV tuner right on the card. This feature lets you display computer data on a standard television without the need for an external scan converter (usually 100 US\$ or more). Business users can then give PC-based presentations with TVs as large-screen monitors, and home users can play computer games on their TV sets. However this feature is also useful for VR users: thanks to the converter it is possible to use – without any extra hardware – the new low-cost DVD oriented head-mounted displays from Olympus (EyeTrek, 600 US\$) or Sony (Glasstron PLM-A35, 500 US\$).

On the software side, an interesting low cost solution is the use of 3D engines included in commercial 3D games for developing simple virtual environments. Many 3D games (50 US\$ each), such as Quake or Unreal, include level editors that allow the user to customize the environments and the avatars. Moreover, Discreet has released free software, *gmax*TM, that allows a professional customization of 3D games. Intended to be a fully capable 3D level editing, modeling, animation, and texture-mapping tool, *gmax* ships with a full suite of professional 3D content and animation features. Discreet approved game developers can publish *gmax* “game packs”, which customize the downloadable version of *gmax* into a fully featured level editor for supported game titles. Using this software, it is possible to edit and create 3D environments, materials, 3D objects, weapons, images and lights.

Obviously, level editing does not allow full control of the environment. In particular, the user interaction with the 3D objects

is usually very limited. To overcome this limitation, now there are different VR development toolkits available for PCs, ranging from high-end authoring toolkits that require significant programming experience to simple “hobbyist” packages. Despite the differences in the types of virtual worlds these products can deliver, the various tools are based on the same VR-development model: they allow users to create or import 3D objects, to apply behavioral attributes such as weight and gravity to the objects, and to program the objects to respond to the user via visual and/or audio events. Ranging in prices from free (<http://www.alice.org>) to 5000 US\$ (Virtools Dev 2.5 or Sense 8 WorldUp R5), the toolkits are the most functional of the available VR software options. While some of them rely exclusively on C or C++ programming to build a virtual world, others offer simpler point-and-click operations to develop a simulation. Using VR toolkits, it is also possible to bring in files from a wide array of software packages, such as Wavefront, 3D Studio, EDS Unigraphics, Pro Engineer, and Intergraph EMS, and they can also import VRML and Multigen databases as well as animation scripts and sounds.

5. Challenges and Open Issues

5.1 Technical Challenges

Even if the significant advances in computer and graphic technology drastically improved the characteristics of a typical VE, VR is still limited by the maturity of the systems available. Even today, no off-the-shelf solutions are available. So, the set up of a VR system usually requires a lot of patience for dealing with conflicting hardware or lacking drivers. Nearly every VR system requires a dedicated staff or at least computer technician to keep the system running smoothly. Moreover, much VR technology is still uncomfortable or unpleasant to use. In particular here are listed some current VR technology limitations for users [91]:

- virtual acoustic displays that require a great deal of computational resources in

order to simulate a small number of sources;

- force and tactile displays, still in their infancies, with limited functionality;
- image generators that can't provide low-latency rendering of head tracked complex scenes, requiring severe trade-offs between performance and scene quality;
- position trackers with small working volumes, inadequate robustness, and problems of latency and poor registration.
- HMDs with limited field of view, and encumbering form factor.

As we have seen, a typical area for VR applications is surgery. However, there have been few developments in the area of tactile feedback. The ability to feel tissue is important. Procedures that require palpitation, such as artery localization and tumor detection, are extremely difficult when the only form of haptic exploration is in the form of forces transmitted through long, clumsy instruments. As noted by Moline [92], "The ability to remotely sense small scale shape information and feel forces that mesh with natural hand motions would greatly improve the performance of minimally invasive surgery and bring a greater sense of realism to virtual trainers" (p. 21).

5.2 Safety Issues

The introduction of patients and clinicians to VEs raises particular safety and ethical issues [45]. In fact, despite developments in VR technology, some users still experience health and safety problems associated with VR use [93]. The key concern from the literature is VR-induced sickness, which could lead to problems [94] including:

- symptoms of motion sickness;
- strain on the ocular system;
- degraded limb and postural control;
- reduced sense of presence;
- the development of responses inappropriate for the real world, which might lead to negative training.

The improved quality of the VR systems is drastically reducing the occurrence of sim-

ulation sickness. For instance, a recent review of clinical applications of VR reported instances of simulation sickness are few and nearly all are transient and minor [6]. In general, for a large proportion of VR users these effects are mild and subside quickly [93].

Nonetheless, patients exposed to virtual reality environments may have disabilities that increase their susceptibility to side effects. Precautions should be taken to ensure the safety and well being of patients, including established protocols for monitoring and controlling exposure to virtual reality environments.

Strategies are needed to detect any adverse effects of exposure, some of which may be difficult to anticipate, at an early stage. According to Lewis and Griffin [94] exposure management protocols for patients in virtual environments should include:

- Screening procedures to detect individuals who may present particular risks.
- Procedures for managing patient exposure to VR applications to ensure rapid adaptation with minimum symptoms.
- Procedures for monitoring unexpected side effects and for ensuring that the system meets its design objectives.

Finally, the effect of VEs on cognition is not fully understood. In a recent report, the US National Advisory Mental Health Council [95] suggested that "Research is needed to understand both the positive and the negative effects [of VEs]... on children's and adult's perceptual and cognitive skills". Such research will require the merging of knowledge from a variety of disciplines including (but not limited to) neuropsychology, neuroimaging, educational theory and technology, human factors, medicine, and computer science.

5.3 Research and Clinical Issues

In the last five years there has been a steady growth in the use of virtual reality in health care due to the advances in information technology and to the decline in costs [4]. As we have seen, using the "virtual reality" keyword we can find 951 papers listed in MEDLINE and 708 in PSYCINFO (all

fields query, accessed June 9, 2003). Much of this growth, however, has been in the form of feasibility studies and pilot trials.

The "best" evidence in evaluating the efficacy of a therapy/approach is the results of randomized, controlled clinical trials. However, if we check the available literature we can find only seventeen controlled trials (see Table 2).

Three tested the training possibilities offered by VR: in surgical training and in teaching physical diagnosis skills. Twelve verified the effectiveness of VR in the treatment of four psychological disorders: acrophobia, body image disturbances, binge eating disorders and fear of flying. The final study analyzed the use of VR in the treatment of adult burn pain.

Why there are so few controlled trials in VR research? The possible answers are three.

First, the lack of standardization in VR devices and software. To date, very few of the various VR systems available are interoperable. This makes difficult their use in contexts other than those in which they were developed.

Second, the lack of standardized protocols that can be shared by the community of researchers. If we check the two clinical databases, we can find only four published clinical protocols: for the treatment of eating disorders [96], fear of flying [97], fear of public speaking [98] and panic disorders [99].

Finally, the costs required for the set-up trials. As we have just seen, the lack of interoperable systems added to the lack of clinical protocols force most researchers to spend a lot of time and money in designing and developing their own VR application: many of them can be considered "one-off" creations tied to a proprietary hardware and software, which have been tuned by a process of trial and error. According to the European funded project VEPSY Updated [100], the cost required for designing a clinical VR application from scratch and testing it on clinical patients using controlled trials may range between 150,000 and 200,000 US\$. As noted by a recent report prepared by the US National Research Council [101], "the government support has been the single most important source of

sustained funding for innovative research in both computer graphics and VR. Beginning in the 1960s with its investments in computer modeling, flight simulators, and visualization techniques, and continuing through current developments in virtual worlds, the federal government has made significant investments in military, civilian, and university research that laid the groundwork for one of today's most dynamic technologies. The commercial pay-offs have included numerous companies formed around federally funded research in graphics and VR" (p. 227). In Europe the most important source of funding for health care VR applications was the European Commission through its Information Society Technology programme. However, in the last five years the funds for VR research coming from the European Commission has been between one-third and one-fifth of the total amount distributed by the US government.

6. Conclusions

In general, the review of current applications shows that VR can be considered a useful tool for diagnosis, therapy, education and training. However, several barriers still remain. The PC-based systems, while inexpensive and easy-to-use, still suffer from a lack of flexibility and capabilities necessary to individualize environments for each patient [85]. On the other hand, in most circumstances the clinical skills of the therapist remain the most important factor in the successful use of VR systems. It is clear that building new and additional virtual environments is important so therapists will continue to investigate applying these tools in their day-to-day clinical practice [6]. Further, many of the actual VR applications are in the clinical investigation or laboratory stage, as clearly showed by the lack of controlled trials.

Significant efforts are still required to move VR into commercial success and therefore routine clinical use. Possible future scenarios will involve multi-disciplinary teams of engineers, computer programmers, and therapists working in con-

cert to treat specific clinical problems. Finally, communication networks have the potential to transform VEs into shared worlds in which individuals, objects, and processes interact without regard to their location. In the future, such networks will probably merge VR and telemedicine applications allowing us to use VE for such purposes as distance learning, distributed training, and e-therapy.

It is hoped that by bringing together this community of experts, further stimulation of interest from granting agencies will be accelerated. Information on advances in VR technology must be made available to the health care community in a format that is easy-to-understand and invites participation [102]. Future potential applications of VR are really only limited by the imaginations of talented individuals.

Acknowledgments

The present work was supported by the Commission of the European Communities (CEC), in particular by the IST programme (Project VEPSY UPDATED, IST-2000-25323, <http://www.cybertherapy.info>, <http://www.e-therapy.info>; Project EMMA, IST-2001-39192).

References

- Satava RM, Jones SB. Medical applications of virtual reality. In: Stanney KM, editor. *Handbook of Virtual Environments: Design, Implementation, and Applications*. Mahwah, NJ: Lawrence Erlbaum Associates, Inc.; 2002. pp. 368-391.
- Wootton R. Telemedicine: an introduction. In: Wootton R, editor. *European Telemedicine 1998/99*. London: Kensington Publications Ltd; 1999. pp. 10-12.
- Riva G. From Telehealth to E-health: Internet and distributed virtual reality in health care. *CyberPsychology & Behavior* 2000; 3 (6): 989-98.
- Riva G. Virtual reality for health care: the status of research. *Cyberpsychology & Behavior* 2002; 5 (3): 219-25.
- Chinnock C. Virtual reality in surgery and medicine. *Hosp Technol Ser* 1994; 13 (18): 1-48.
- Riva G, Wiederhold B, Molinari E, editors. *Virtual environments in clinical psychology and neuroscience: Methods and techniques in advanced patient-therapist interaction*. Amsterdam: IOS Press. Online: <http://www.cybertherapy.info/pages/book2.htm>; 1998.
- Beolchi L, Riva G. Virtual reality for health care. In: Akay M, Marsh A, editors. *Information Technologies in Medicine*. Toronto: John Wiley & Sons; 2001. pp. 39-83.
- Riva G, Wiederhold BK. Introduction to the special issue on virtual reality environments in behavioral sciences. *IEEE Transactions on Information Technology in Biomedicine* 2002; 6 (3): 193-7.
- Rubino F, Soler L, Marescaux J, Maisonneuve H. Advances in virtual reality are wide ranging. *Bmj* 2002; 324 (7337): 612.
- McCloy R, Stone R. Science, medicine, and the future. Virtual reality in surgery. *Bmj* 2001; 323 (7318): 912-5.
- Székelly G, Satava RM. Virtual reality in medicine. *Bmj* 1999; 319 (7220): 1305.
- Schultheis MT, Rizzo AA. The Application of Virtual Reality Technology in Rehabilitation. *Rehabilitation Psychology* 2001; 46 (3): 296-311.
- Riva G, Rizzo A, Alpini D, Attree EA, Barbieri E, Bertella L, et al. Virtual environments in the diagnosis, prevention, and intervention of age-related diseases: A review of VR scenarios proposed in the EC VETERAN project. *CyberPsychology and Behavior* 1999; 2 (6): 577-91.
- Rizzo AA, Wiederhold B, Riva G, Van Der Zaag C. A bibliography of articles relevant to the application of virtual reality in the mental health field. *CyberPsychology & Behavior* 1998; 1 (4): 411-25.
- Rizzo AA, Buckwalter JG. Virtual reality and cognitive assessment and rehabilitation: the state of the art. In: Riva G, editor. *Virtual reality in neuro-psycho-physiology*. Amsterdam: IOS Press; 1997. pp. 123-146. Online: <http://www.cybertherapy.info/pages/book1.htm>.
- Riva G, Alcañiz M, Anolli L, Bacchetta M, Baños RM, Beltrame F, et al. The VEPSY Updated project: Virtual reality in clinical psychology. *CyberPsychology and Behavior* 2001; 4 (4): 449-55.
- Heim M. *Virtual Realism*. New York: Oxford University Press; 1998.
- Bricken W. *Virtual reality: Directions of growth*. Seattle, WA: University of Washington; 1990. Report No.: HITL Technical Report R-90-1.
- Riva G, Mantovani G. The need for a socio-cultural perspective in the implementation of virtual environments. *Virtual Reality* 2000 (5): 32-8.
- Riva G, Davide F, editors. *Communications through Virtual Technologies: Identity, Community and Technology in the Communication Age*. Amsterdam: IOS Press. Online: <http://www.emergingcommunication.com/volume1.html>; 2001.
- Biocca F, Levy MR, editors. *Communication in the age of virtual reality*. Hillsdale, NJ: Lawrence Erlbaum Associates; 1995.
- Steuer JS. Defining virtual reality: Dimensions determining telepresence. *Journal of Communication* 1992; 42 (4): 73-93.
- Sastry L, Boyd DRS. Virtual environments for engineering applications. *Virtual Reality: Research, development and applications* 1998; 3 (4): 235-44.
- Satava RM, Ellis SR. Human interface technology. An essential tool for the modern surgeon. *Surg Endosc* 1994; 8 (7): 817-20.

25. Rizzo AA, Neumann U, Enciso R, Fidaleo D, Noh JY. Performance-driven facial animation: basic research on human judgments of emotional state in facial avatars. *Cyberpsychol Behav* 2001; 4 (4): 471-87.
26. Riva G, Davide F, IJsselstein WA, editors. *Being There: Concepts, effects and measurements of user presence in synthetic environments*. Amsterdam: IOS Press. Online: <http://www.emergingcommunication.com/volume5.html>; 2003.
27. IJsselstein WA, Lombard M, Freeman J. Toward a core bibliography of presence. *Cyberpsychology & Behavior* 2001; 4 (2): 317-21.
28. Riva G, Waterworth JA. Presence and the Self: A cognitive neuroscience approach. *Presence-Connect* 2003; 3 (1): Online: <http://presence.cs.ucl.ac.uk/presenceconnect/articles/Apr2003/jwworthApr72003114532/jwworthApr72003114532.html>.
29. Slater M, Wilbur S. A framework for immersive virtual environments (FIVE): Speculations on the role of presence in virtual environments. *Presence: Teleoperators and Virtual Environments* 1997; 6 (6): 603-16.
30. Waterworth JA, Waterworth EL. Focus, Locus, and Sensus: The three dimensions of virtual experience. *Cyberpsychology and Behavior* 2001; 4 (2): 203-13.
31. Mantovani G, Riva G. "Real" presence: How different ontologies generate different criteria for presence, telepresence, and virtual presence. *Presence, Teleoperators, and Virtual Environments* 1999; 8 (5): 538-48.
32. Schubert T, Friedman F, Regenbrecht H. The experience of presence: Factor analytic insights. *Presence: Teleoperators, and Virtual Environments* 2001; 10 (3): 266-81.
33. Zahoric P, Jenison RL. Presence as being-in-the-world. *Presence, Teleoperators, and Virtual Environments* 1998; 7 (1): 78-89.
34. Janoff-Bulman R, Frantz CM. The impact of trauma on meaning: From meaningless world to meaningful life. In: Power M, Brewin CR, editors. *The transformation of meaning in psychological therapies*. New York: Wiley; 1997. pp. 91-106.
35. Damasio A. *The Feeling of What Happens: Body, Emotion and the Making of Consciousness*. San Diego, CA: Harcourt Brace and Co, Inc.; 1999.
36. ISO. ISO/IEC 9126-Software engineering – Product quality – Part 1: Quality model. Geneva: International Organization for Standardization; 2001.
37. John NW, Thacker N, Pokric M, Jackson A, Zanetti G, Gobetti E, et al. An integrated simulator for surgery of the petrous bone. *Stud Health Technol Inform* 2001; 81: 218-24.
38. Dobson HD, Pearl RK, Orsay CP, Rasmussen M, Evenhouse R, Ai Z, et al. Virtual reality: new method of teaching anorectal and pelvic floor anatomy. *Dis Colon Rectum* 2003; 46 (3): 349-52.
39. Alcañiz M, Perpiña C, Baños R, Lozano JA, Montesa J, Botella C, et al. A new realistic 3D body representation in virtual environments for the treatment of disturbed body image in eating disorders. *CyberPsychology and Behavior* 2000; 3 (3): 421-32.
40. Ackerman MJ. The Visible Human Project. *J Biocommun* 1991; 18 (2): 14.
41. Spitzer V, Ackerman MJ, Scherzinger AL, Whitlock D. The visible human male: a technical report. *J Am Med Inform Assoc* 1996; 3 (2): 118-30.
42. Ackerman MJ, Yoo T, Jenkins D. From data to knowledge – the Visible Human Project continues. *Medinfo* 2001; 10 (Pt 2): 887-90.
43. Westwood JD, Hoffman HM, Mogel GT, Stredney D, editors. *Medicine meets virtual reality 2002*. Amsterdam: IOS Press; 2002.
44. Jeffries PR, Woolf S, Linde B. Technology-based vs. traditional instruction. A comparison of two methods for teaching the skill of performing a 12-lead ECG. *Nurs Educ Perspect* 2003; 24 (2): 70-4.
45. Durlach NI, Mavor ASE. *Virtual reality: scientific and technological challenges*. Washington, D.C.: National Academy Press. Online: <http://www.nap.edu/books/0309051355/html/index.html>; 1995.
46. Satava RM. *Surgery 2001: A Technologic Framework for the Future*. *Surgical Endoscopy* 1993; 7: 111-3.
47. Satava RM. Virtual reality surgical simulator. The first steps. *Surg Endosc* 1993; 7 (3): 203-5.
48. Satava RM. Surgical education and surgical simulation. *World J Surg* 2001; 25 (11): 1484-9.
49. Sung WH, Fung CP, Chen AC, Yuan CC, Ng HT, Doong JL. The assessment of stability and reliability of a virtual reality-based laparoscopic gynecology simulation system. *Eur J Gynaecol Oncol* 2003; 24 (2): 143-6.
50. Friedl R, Preisack MB, Klas W, Rose T, Stracke S, Quast KJ, et al. Virtual Reality and 3D Visualizations in Heart Surgery Education. *Heart Surg Forum* 2002; 5 (3): E17-21.
51. Ali MR, Mowery Y, Kaplan B, DeMaria EJ. Training the novice in laparoscopy. *Surg Endosc* 2002.
52. Gor M, McCloy R, Stone R, Smith A. Virtual reality laparoscopic simulator for assessment in gynaecology. *Bjog* 2003; 110 (2): 181-7.
53. Dammann F, Bode A, Schwaderer E, Schleich M, Heuschmid M, Maassen MM. Computer-aided surgical planning for implantation of hearing aids based on CT data in a VR environment. *Radiographics* 2001; 21 (1): 183-91.
54. Xia J, Ip HH, Samman N, Wong HT, Gateno J, Wang D, et al. Three-dimensional virtual-reality surgical planning and soft-tissue prediction for orthognathic surgery. *IEEE Trans Inf Technol Biomed* 2001; 5 (2): 97-107.
55. Herfarth C, Lamade W, Fischer L, Chiu P, Cardenas C, Thorn M, et al. The effect of virtual reality and training on liver operation planning. *Swiss Surg* 2002; 8 (2): 67-73.
56. Ross MD, Twombly IA, Bruyns C, Cheng R, Senger S. Telecommunications for health care over distance: the virtual collaborative clinic. *Studies in Health Technology and Informatics* 2000; 70: 286-91.
57. Erel E, Aiyenibe B, Butler PE. Microsurgery simulators in virtual reality: Review. *Microsurgery* 2003; 23 (2): 147-52.
58. Halligan S, Fenlon HM. Virtual colonoscopy. *Bmj* 1999; 319 (7219): 1249-52.
59. Moorthy K, Smith S, Brown T, Bann S, Darzi A. Evaluation of virtual reality bronchoscopy as a learning and assessment tool. *Respiration* 2003; 70 (2): 195-9.
60. Dunkin BJ. Flexible endoscopy simulators. *Semin Laparosc Surg* 2003; 10 (1): 29-35.
61. Vincelli F. From imagination to virtual reality: the future of clinical psychology. *CyberPsychology & Behavior* 1999; 2 (3): 241-8.
62. Vincelli F, Molinari E, Riva G. Virtual reality as clinical tool: immersion and three-dimensionality in the relationship between patient and therapist. *Studies in Health Technology and Informatics* 2001; 81: 551-3.
63. Norcross JC, Hedges M, Prochaska JO. The face of 2010: A Delphi poll on the future of psychotherapy. *Professional Psychology: Research and Practice* 2002; 33 (3): 316-22.
64. Menon ST. Psychological Empowerment: Definition, Measurement, and Validation. *Canadian Journal of Behavioural Science* 1999; 31 (3): 161-4.
65. Botella C, Perpiña C, Baños RM, Garcia-Palacios A. Virtual reality: a new clinical setting lab. *Studies in Health Technology and Informatics* 1998; 58: 73-81.
66. Emmelkamp PM, Bruynzeel M, Drost L, van der Mast CA. Virtual reality treatment in acrophobia: a comparison with exposure in vivo. *CyberPsychology & Behavior* 2001; 4 (3): 335-9.
67. Emmelkamp PM, Krijn M, Hulsbosch AM, de Vries S, Schuemie MJ, van der Mast CA. Virtual reality treatment versus exposure in vivo: a comparative evaluation in acrophobia. *Behaviour Research & Therapy* 2002; 40 (5): 509-16.
68. Rothbaum BO, Hodges LF, Kooper R, Opdyke D, Williford JS, North M. Effectiveness of computer-generated (virtual reality) graded exposure in the treatment of acrophobia. *Am J Psychiatry* 1995; 152 (4): 626-8.
69. Garcia-Palacios A, Hoffman H, Carlin A, Furness TA, 3rd, Botella C. Virtual reality in the treatment of spider phobia: a controlled study. *Behavior Research and Therapy* 2002; 40 (9): 983-93.
70. Vincelli F, Anolli L, Bouchard S, Wiederhold BK, Zurloni V, Riva G. Experiential Cognitive Therapy in the treatment of Panic Disorders with Agoraphobia: A controlled study. *CyberPsychology & Behavior* 2003; 6 (3): 312-8.
71. Riva G, Bacchetta M, Baruffi M, Molinari E. Virtual reality-based multidimensional therapy for the treatment of body image disturbances in obesity: a controlled study. *Cyberpsychology and Behavior* 2001; 4 (4): 511-26.
72. Riva G, Bacchetta M, Baruffi M, Molinari E. Virtual-reality-based multidimensional therapy for the treatment of body image disturbances in binge eating disorders: a preliminary controlled study. *IEEE Transactions on Information Technology in Biomedicine* 2002; 6 (3): 224-34.
73. Riva G, Bacchetta M, Cesa G, Conti S, Molinari E. Six-month follow-up of in-patient Expe-

- ritional-Cognitive Therapy for binge eating disorders. *CyberPsychology & Behavior* 2003; 6 (3): 251-8.
74. Rothbaum BO, Hodges L, Smith S, Lee JH, Price L. A controlled study of virtual reality exposure therapy for the fear of flying. *Journal of Consulting & Clinical Psychology* 2000; 68 (6): 1020-6.
 75. Rothbaum BO, Hodges L, Anderson PL, Price L, Smith S. Twelve-month follow-up of virtual reality and standard exposure therapies for the fear of flying. *J Consult Clin Psychol* 2002; 70 (2): 428-32.
 76. Maltby N, Kirsch I, Mayers M, Allen G. Virtual Reality Exposure Therapy for the treatment of fear of flying: A controlled investigation. *Journal of Consulting & Clinical Psychology* 2002; 70 (5): 1112-8.
 77. Wiederhold BK, Jang DP, Kim SI, Wiederhold MD. Physiological monitoring as an objective tool in virtual reality therapy. *Cyberpsychology & Behavior* 2002; 5 (1): 77-82.
 78. Wiederhold BK, Jang DP, Gevirtz RG, Kim SI, Kim IY, Wiederhold MD. The treatment of fear of flying: a controlled study of imaginal and virtual reality graded exposure therapy. *IEEE Transactions on Information Technology in Biomedicine* 2002; 6 (3): 218-23.
 79. Riva G. Virtual environments in neuroscience. *IEEE Transactions on Information Technology in Biomedicine* 1998; 2 (4): 275-81.
 80. Rose FD, Brooks BM, Attree EA, Parslow DM, Leadbetter AG, McNeil JE, et al. A preliminary investigation into the use of virtual environments in memory retraining after vascular brain injury: indications for future strategy? *Disabil Rehabil* 1999; 21 (12): 548-54.
 81. Gourlay D, Lun KC, Lee YN, Tay J. Virtual reality for relearning daily living skills. *Int J Med Inf* 2000; 60 (3): 255-61.
 82. Riva G. Virtual reality in rehabilitation of spinal cord injuries. *Rehabilitation Psychology* 2000; 45 (1): 81-8.
 83. Sisto SA, Forrest GF, Glendinning D. Virtual reality applications for motor rehabilitation after stroke. *Topics in Stroke Rehabilitation* 2002; 8 (4): 11-23.
 84. Tarr MJ, Warren WH. Virtual reality in behavioral neuroscience and beyond. *Nature Neuroscience* 2002; 5 Suppl: 1089-92.
 85. Riva G, editor. Virtual reality in neuro-psycho-physiology: Cognitive, clinical and methodological issues in assessment and rehabilitation. Amsterdam: IOS Press. Online: <http://www.cybertherapy.info/pages/book1.htm>; 1997.
 86. Zhang L, Abreu BC, Masel B, Scheibel RS, Christiansen CH, Huddleston N, et al. Virtual reality in the assessment of selected cognitive function after brain injury. *Am J Phys Med Rehabil* 2001; 80 (8): 597-604; quiz 605.
 87. Wald J, Liu L. Psychometric properties of the driVR: a virtual reality driving assessment. *Studies in Health Technology and Informatics* 2001; 81: 564-6.
 88. Pugnelli L, Mendozzi L, Motta A, Cattaneo A, Barbieri E, Brancotti A. Evaluation and re-training of adults' cognitive impairment: which role for virtual reality technology? *Comput Biol Med* 1995; 25 (2): 213-27.
 89. Broeren J, Bjorkdahl A, Pascher R, Rydmark M. Virtual reality and haptics as an assessment device in the postacute phase after stroke. *Cyberpsychol Behav* 2002; 5 (3): 207-11.
 90. Piron L, Cenni F, Tonin P, Dam M. Virtual Reality as an assessment tool for arm motor deficits after brain lesions. *Stud Health Technol Inform* 2001; 81: 386-92.
 91. Gross D. Technology Management and User Acceptance of VE Technology. In: Stanney KM, editor. *Handbook of Virtual Environments: Design, Implementation, and Applications*. Mahwah, NJ: Lawrence Erlbaum Associates, Inc.; 2002.
 92. Moline J. Virtual reality in health care: a survey. In: Riva G, editor. *Virtual reality in neuro-psycho-physiology*. Amsterdam: IOS Press; 1997: pp. 3-34.
 93. Nichols S, Patel H. Health and safety implications of virtual reality: a review of empirical evidence. *Appl Ergon* 2002; 33 (3): 251-71.
 94. Lewis CH, Griffin MJ. Human factors consideration in clinical applications of virtual reality. In: Riva G, editor. *Virtual reality in neuro-psycho-physiology*. Amsterdam: IOS Press; 1997: pp. 35-56.
 95. A report of the U.S. National Advisory Mental Health Council. NIH Publication. Washington, D.C.: U.S. Government Printing Office; 1995. Report No.: 95-3682.
 96. Riva G, Bacchetta M, Cesa G, Conti S, Molinari E. Virtual reality and telemedicine based Experiential Cognitive Therapy: Rationale and Clinical Protocol. In: Riva G, Galimberti C, editors. *Towards CyberPsychology: Mind, Cognition and Society in the Internet Age*. Amsterdam: IOS Press; 2001: pp. 273-308.
 97. Klein RA. Treating fear of flying with virtual reality exposure therapy. In: VandeCreek, Leon (Ed); Jackson, Thomas L. (Ed). (1999). *Innovations in clinical practice: A source book*, Vol. 17. (pp. 449-465). Sarasota, FL, US.; 1999.
 98. Botella C, Baños RM, Villa H, Perpiña C, Garcia-Palacios A. Telepsychology: Public speaking fear treatment on the internet. *CyberPsychology and Behavior* 2000; 3 (6): 959-68.
 99. Vincelli F, Choi YH, Molinari E, Wiederhold BK, Riva G. A VR-based multicomponent treatment for panic disorders with agoraphobia. *Studies in Health Technology and Informatics* 2001; 81: 544-50.
 100. Riva G, Bolzoni M, Carella F, Galimberti C, Griffin MJ, Lewis CH, et al. Virtual reality environments for psycho-neuro-physiological assessment and rehabilitation. In: Morgan KS, Weghorst SJ, Hoffman HM, Stredney D, editors. *Medicine Meets Virtual Reality: Global Healthcare Grid*. Amsterdam: IOS Press; 1997: pp. 34-45.
 101. Hughes T, Clark DD, Banks PM, Lineberger WC, editors. *Funding a Revolution: Government Support for Computing Research*. Washington, DC: National Academy Press. Online: <http://stills.nap.edu/html/far/contents.html>; 1999.
 102. Riva G, Bacchetta M, Baruffi M, Borgomainerio E, Defrance C, Gatti F, et al. VREPAR Projects: The use of virtual environments in psycho-neuro-physiological assessment and rehabilitation. *CyberPsychology & Behavior* 1999; 2 (1): 69-76.

Correspondence to:
 Prof. Giuseppe Riva, Ph.D.
 Dipartimento di Psicologia
 Università Cattolica del Sacro Cuore
 Largo Gemelli 1
 20123, Milan, Italy
 E-mail: auxo.psylab@auxologico.it
 Web site: <http://www.atmplab.com>